

Pediatric Palliative Care Needs in the Canton of Bern, Switzerland

A Cross-Sectional Study

Brigitte Hofer Fritzsche, Stiftung Kifa; Eva Maria Tinner Oehler, Kinderspitäler Zürich und Bern; Maya Zumstein-Shaha, Berner Fachhochschule, Schweiz.

Challenges




- In Switzerland, 450 – 500 children die each year
 - The needs of children in Paediatric Palliative Care (PPC) are different compared to the needs of adult patients with palliative care:
 - ✓ The time PPC is required varies from less than one month up to several years
 - ✓ Early inclusion of PPC in the disease trajectory, i.e., concurrent to curative treatment, is strongly recommended
 - ✓ Support of the child's family is an essential part of PPC
 - ✓ Adjusted support is necessary for promoting physiological child development
 - Today, no comprehensive care network for PPC exists in the Canton of Bern
 - Prevalence data on the number of children needing PPC does not exist

Aims



- To determine the number of children with life-limiting conditions living in the Canton of Bern who are cared for by Home Healthcare Organisations and who meet the criteria for PPC according to the START-IMPACT screening tool (version 1.8.)
 - To describe the characteristics of the affected children and their families

Methods



Design: Sample:


Cross-sectional study, patient dossier analysis

Recruitment:


Complete survey of all registered children due to the heterogeneity of the study-population and the estimated low number of children with life-limiting conditions in the study area

Recruitment:

and the estimated low number of children with life limiting conditions in the study area
Children cared for by Home Healthcare, age 0 -20 years, with a START-IMPACT Score ≥ 4



Discussion



- The burden of illness and care of the affected children and their families is enormous
 - Access to specialized PPC Services is inadequate
 - ✓ Early inclusion of PPC in the disease trajectory is strongly recommended
 - Care and treatment gaps in various areas exist
 - ✓ High care and treatment burden of the families
 - ✓ Elevated risk for depression, sleeping problems, social Isolation and financial problems → mothers are more often affected
 - Establishment of an adequate comprehensive care network for PPC in Bern needed
 - ✓ ACP and the continuous care by an interprofessional specialized PPC team (in- and outpatient) is recommended

